

Published online:

29 June 2024

 Web version

# TWA1: a novel thermosensor enhancing plant thermotolerance

by Mauro Maver 

**In this study, Bohn et al. identified THERMO-WITH-ABA-RESPONSE 1 as a novel thermosensor in *Arabidopsis thaliana*. This sensor is crucial for thermotolerance and heat-stress response as it interacts with key transcription factors and co-repressors in a temperature-dependent mechanism. This discovery has the potential to significantly improve crop acclimation to global warming through biotechnological applications.**

In the plant kingdom, each species has an optimal temperature range for growth and development. However, sudden climatic fluctuations or prolonged periods of intense heat can severely challenge their survival. When temperatures exceed this range, plants activate heat-stress responses (HSR) to adapt to these stressful conditions. These adaptation and response strategies involve various developmental processes in the plant<sup>1</sup>. Heat triggers the HSR through heat-shock factor 1 (HSF1) transcription factors, which induce the expression of heat-shock proteins (HSPs), helping the plant adapt to heat. Additionally, the HSR involves several phytohormone pathways, with abscisic acid (ABA) playing a crucial role in regulating the plant's water status, promoting thermotolerance, and enhancing plant acclimation<sup>2,3</sup>.

To better understand the mechanisms of temperature perception and the induction of thermotolerance, Bohn et al. examined different lines of *Arabidopsis thaliana* that were hyper-responsive or insensitive to ABA under heat stress. Among all the mutants tested, one showed strong thermosensitivity, and the gene locus was subsequently named *THERMO-WITH-ABA-RESPONSE 1* (*TWA1*). Expressed throughout the plant, it encodes a predicted intrinsically disordered protein of 130 kDa with two potential ethylene-responsive-element-binding-factor-associated amphiphilic repression (EAR) motifs (LxLxL).

In a yeast two-hybrid (Y2H) system, *TWA1* was found to interact with

JASMONATE-ASSOCIATED MYC-LIKE 2 (*JAM2*), a transcription factor involved in jasmonate (JA) hormone signalling<sup>4,5</sup>, and with the co-repressors *TOPLESS* (*TPL*) and *TOPLESS-RELATED* (*TPR*), forming a repressor complex. However, this interaction is highly temperature-dependent. Specifically, *TWA1* only accumulates in nuclear subdomains at elevated temperatures (30–35°C), and its conformational changes allow physical interaction with *JAM2* and *TPL/TPR*. These temperature-dependent structural rearrangements are mediated by a 20 amino acid sequence in the amino-terminal region of *TWA1*, defined as the Highly Variable Region (HVR). The HVR thus confers strong thermosensitivity to *TWA1*.

Homologues of *TWA1* have been identified in both monocotyledons and dicotyledons. An interesting aspect is the role of HVR in thermal sensing, particularly its amino acid sequence. Although they have comparable inhibitory effects at 30–35°C, their thermal sensitivity or temperature threshold varies. For instance, half-maximum inhibitory values ( $IT_{50}$ ) of 20°C, 26°C, and 30°C were observed for *TWA1* in *A.lyrata* (cold acclimated), *A.thaliana*, and *Sinapis alba* (Mediterranean climate), respectively. Similarly, this threshold has been shown to affect accumulation and interaction with *JAM2* and *TPL* and, consequently, the activation of its inhibitor effects.

Moreover, tests involving the overexpression of *TWA1* (*TWA1*(oe)) have demonstrated its ability to confer an enhanced thermotolerance, both basal and acquired, without impacting growth and photosynthesis. No significant differences in growth, photosynthesis, and gas exchange were observed between WT and *TWA1*(oe) when grown at 20°C. These findings reconfirm the activation of *TWA1* only above specific temperatures. Additionally, the improved thermotolerance of *TWA1* is not due to the constitutive upregulation of HSP transcripts.

*TWA1* presents an interesting parallel with other known thermal sensors, such as *EARLY FLOWERING 3* (*ELF3*) in *A.thaliana*<sup>6</sup>. However, thermally, *TWA1* acts oppositely to *ELF3*, repressing gene expression at high temperatures and being much more sensitive to temperature

changes.

In conclusion, Bohn et al. have identified a new thermosensor, *TWA1*, and demonstrated its requirement for thermotolerance and the transcriptional upregulation of *HSFA2* and *HSPs* in *A.thaliana*. The potential of *TWA1* in breeding and biotechnology is vast, providing valuable tools for enhancing crop acclimation to the growing issue of global warming. Additionally, *TWA1* and its orthologues represent a valuable resource in the emerging field of thermogenetics<sup>7</sup>.

HSR, heat-stress responses • HSPs, heat-shock proteins • ABA, abscisic acid • *TWA1*, *THERMO-WITH-ABA-RESPONSE 1* • Da, Dalton • EAR, ethylene-responsive-element-binding-factor-associated amphiphilic repression • Y2H, yeast-to-hybrid • *JAM2*, JASMONATE-ASSOCIATED MYC2-LIKE 2 • *TPL*, *TOPLESS* • *TPR*, *TOPLESS-RELATED* • HVR, highly variable region • *TWA1*(oe), *TWA1*-overexpressed • *ELF3*, *EARLY FLOWERING 3* • *HSFA2*, heat-shock transcription factor A2.

## References

- Casal, J.J. & Balasubramanian S. Thermomorphogenesis. *Ann. Rev. Plant Biol.* **70**, 321–446 (2019).
- Cutler, S.R. et al. Abscisic acid: Emergence of a core signaling network. *Ann. Rev. Plant Biol.* **61**, 651–679 (2010).
- Yoshida, T. et al. Revisiting the basal role of ABA—roles outside of stress. *Trends. Plant Sci.* **24**, 625–635 (2019).
- Sasaki-Sekimoto, Y. et al. helix-loop-helix transcription factors JASMONATE-ASSOCIATED MYC2-LIKE1 (*JAM1*), *JAM2*, and *JAM3* are negative regulators of jasmonate responses in arabidopsis. *Plant Physiol.* **163**, 291–304 (2013).
- Song, S. et al. The bHLH subgroup IIIId factors negatively regulate jasmonate-mediated plant defense and development. *PLoS Genet.* **9**, e1003653 (2013).
- Jung, J.-H. et al. A prion-like domain in *ELF3* functions as a thermosensor in arabidopsis. *Nature* **585**, 256–260 (2020).
- Chee, W.K.D. et al. Thermogenetics: Applications come of age. *Biotechnol. Adv.* **55**, 107907 (2022).

**Original article:** Bohn L. et al. The temperature sensor *TWA1* is required for thermotolerance in *Arabidopsis*. *Nature* **629**, 1126–1132 doi.org/10.1038/s41586-024-07424-x (2024)